

Native Borrelia Burgdorferi Cells

Cat. No. BOR-026 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Heat-killed *Borrelia burgdorferi* cells in dextran solution. Antigen is intended for use as a positive control in immunoassay development for *Borrelia* detection.

Background

Lyme disease is a multisystem disorder caused by three species of the tick-borne spirochete *Borrelia burgdorferi* sensu lato; namely *Borrelia burgdorferi* sensu stricto (United States and Europe), *Borrelia garinii* and *Borrelia afzelii* (Europe). The disease was first recognized in Lyme, Connecticut, United States in 1975, and *Borrelia burgdorferi* was described as the causative organism in 1982. However, the clinical manifestations of the disorder were previously described by Afzelius in Europe in the early 1900's (Zouyan & Coburn, 2017).

Borrelia are loosely coiled, spiral, microaerophilic bacteria with 3 to 10 loose coils arranged in a helical shape. The cells contain at least seven periplasmic flagella that are responsible for the organism's motility (Wormser, 2012). *B. burgdorferi* resembles other spirochetes in that it has an outer membrane and inner membrane with a thin layer of peptidoglycan in between. However, the outer membrane lacks lipopolysaccharide. Its shape is a flat wave. It is about 0.3 µm wide and 5 to 20 µm in length. It is a slow-growing microaerophilic spirochete with a doubling time of 24 to 48 hours. *Borrelia burgdorferi* sensu lato (or *B. burgdorferi* in the general sense) consists of 14 closely related genospecies. However, the human infection is caused primarily by three genospecies. In the United States, *B. burgdorferi* sensu stricto (*B. burgdorferi* in the strict sense), is the sole cause of the disease, whereas *Borrelia afzelii* and *Borrelia garinii* are the primary causes in Europe and Asia (Steere, 2015). Each of these genospecies is associated with different enzootic lifecycles and clinical manifestations. *Borrelia afzelii* is mostly associated with skin manifestations, whereas

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

B. garinii is considered to be the most neurotropic and B. burgdorferi s.s. the most arthritogenic. These genospecies are associated with different vertebrate host species. Borrelia afzelii is associated with rodents, whereas B. garinii is associated with birds (van Duijvendijk et al., 2015).

Notes	This product is intended for research and manufacturing uses only. It is not a diagnostic device. The user assumes all responsibility for care, custody and control of the material, including its disposal, in accordance with all regulations.
--------------	--

Type	Native
-------------	--------

ClassID 1	Infectious Disease
------------------	--------------------

GENE INFORMATION

Synonyms	Borrelia Burgdorferi
-----------------	----------------------

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA