

Recombinant *Borrelia burgdorferi* OspA Protein, His-tagged

Cat. No. BOR-032 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This is a recombinant <i>Borrelia burgdorferi</i> OspA protein, fused to a His-tag and produced in <i>E. coli</i> .
Tag	His
Background	Outer-Surface Protein A (OspA) is a 31 kDa lipoprotein encoded by <i>Borrelia burgdorferi</i> and is a major component of the spirochete's extracellular matrix (Stevenson, et al., 1996), probably associated with lipid rafts and serving as a lipid-anchor (Toledo, et al., 2014). Strain B31 is the type strain (ATCC 35210) for this organism and was derived by limited dilutional cloning from the original Lyme-disease tick isolate obtained by A. Barbour (Johnson, et al., 1984). The <i>Borrelia</i> species causing Lyme disease express different OspA serotypes on their surface, <i>B. burgdorferi</i> (serotype 1), <i>B. afzelii</i> (serotype 2), <i>B. garinii</i> (serotypes, 3, 5 and 6) and <i>B. bavariensis</i> (serotype 4) (Wilske, et al., 1988). Many of the borrelial surface antigens are lipid-modified proteins (i.e. lipoproteins), although a number of these surface-exposed lipoproteins (OspA, OspB, and OspC) are not found exclusively on the surface of the organism. These lipoproteins are also detected in the periplasm of the organism and can be shuttled to and from the borrelial surface at different points during infection. The interface between <i>B. burgdorferi</i> and its human host is its outer surface (and therefore proteins localized to the outer membrane) play an important role in dissemination, virulence, tissue tropism, and immune evasion. Antibodies directed against outer surface proteins have also been shown to protect animals and humans from infection with <i>B. burgdorferi</i> (Kenedy, et al., 2012).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

OspA likely mediates the attachment of *B. burgdorferi* to the tick mid-gut by binding the mid-gut receptor TROSPA (Tick Receptor for OspA). TROSPA is downregulated to allow migration out of the tick mid-gut during feeding, and into the salivary glands before being transmitted to the mammalian host. This transition is believed to be facilitated by changes in expression of some *B. burgdorferi* genes, including OspA (Ding, et al., 2000). These changes may be regulated by changes in tick life cycle, changes in conditions during tick feeding (such as temperature, pH and nutrients) and/or in coordination with the course of infection in the mammal host (Norris, 2006). A heterodimer of the linked C-terminal half of two OspA serotypes was shown to protect mice from a challenge with spirochetes expressing either OspA serotype 1, 2 or 5, when challenged with infected ticks and in vitro grown spirochetes (Comstedt, et al., 2014). OspA-specific human Mabs can prevent the transmission of *B. burgdorferi* from ticks to mice (Wang, et al., 2016). More recently, structural analysis of the human antibody LA-2/OspA complex has revealed specific residues that may be exploited to modulate recognition of the protective epitope of OspA potentially offering a new path towards prophylactic passive antibodies (Shivender, et al., 2017).

Purity	>95%
Formulation	DPBS
Notes	This product is intended for research and manufacturing uses only. It is not a diagnostic device. The user assumes all responsibility for care, custody and control of the material, including its disposal, in accordance with all regulations.
Type	Recombinant
ClassID 1	Infectious Disease

GENE INFORMATION

Synonyms

Borrelia burgdorferi OspA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA