

Recombinant Human TNFSF10 293 Cell Lysate

Cat. No. TNFSF10-892HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

TNFSF10 tumor necrosis factor (ligand) superfamily, member 10 [Homo sapiens]

Official Symbol

TNFSF10

Synonyms

TNFSF10; tumor necrosis factor (ligand) superfamily, member 10; tumor necrosis factor ligand superfamily member 10; Apo 2L; CD253; TL2; TRAIL; Apo-2 ligand; TNF-related apoptosis inducing ligand TRAIL; tumor necrosis factor (ligand) family, member 10; chemokine tumor necrosis factor ligand superfamily member 10; tumor necrosis factor apoptosis-inducing ligand splice variant delta; APO2L; Apo-2L;

Gene ID

8743

mRNA Refseq

NM_001190942

Protein Refseq

NP_001177871

MIM

603598

UniProt ID

P50591

**Chromosome
Location**

3q26

Pathway

Activation of Pro-Caspase 8, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptosis, conserved

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Apoptosis, organism-specific biosystem; Caspase-8 is formed from procaspase-8, organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA