

Recombinant Human pericentrin Protein, His tagged

Cat. No. PCNT-045H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human PCNT protein (O95613) (His 258-Leu 553) with a His tag at the C-terminus was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	258-553aa
Description	Pericentrin (kendrin), also known as PCNT and pericentrin-B (PCNTB), is a protein which in humans is encoded by the PCNT gene on chromosome 21. This protein localizes to the centrosome and recruits proteins to the pericentriolar matrix (PCM) to ensure proper centrosome and mitotic spindle formation, and thus, uninterrupted cell cycle progression. This gene is implicated in many diseases and disorders, including congenital disorders such as microcephalic osteodysplastic primordial dwarfism type II (MOPDII) and Seckel syndrome. The protein encoded by this gene is expressed in the cytoplasm and centrosome throughout the cell cycle, and to a lesser extent, in the nucleus. It is an integral component of the PCM, which is a centrosome scaffold that anchors microtubule nucleating complexes and other centrosomal proteins. In one model, PCNT complexes with CEP215 and is phosphorylated by PLK1, leading to PCM component recruitment and organization, centrosome maturation, and spindle formation. The protein controls the nucleation of microtubules by interacting with the microtubule nucleation component γ-tubulin, thus anchoring the γ-tubulin ring complex to the centrosome, which is essential for bipolar spindle formation and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

chromosome assembly in early mitosis. This ensures normal function and organization of the centrosomes, mitotic spindles, and cytoskeleton, and by extension, regulation over cell cycle progression and checkpoints.

Tag	C-His
Molecular Mass	36.7 kDa
Endotoxin	Less than 1.0 EU/μg by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade. It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage Buffer	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.
Reconstitution	Reconstitute at 250 μg/mL in sterile water.

GENE INFORMATION

Gene Name	PCNT pericentrin [Homo sapiens (human)]
Official Symbol	PCNT
Synonyms	PCNT; pericentrin; PCNT2, pericentrin 2 (kendrin); KEN; kendrin; KIAA0402; PCN; PCNTB; SCKL4; Seckel syndrome 4; pericentrin B; pericentrin-2; pericentrin-B; pericentrin-380; MOPD2; PCNT2; PCTN2;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 5116


mRNA Refseq NM_006031

Protein Refseq NP_006022

MIM 605925

UniProt ID O95613

**Protein on SDS-
PAGE under
reducing (R)
condition.**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA