

Active Recombinant Full Length Human deoxycytidine kinase Protein, R104M, D133A, His tagged

Cat. No. DCK-05HFL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant full length (aa 1-260) wild-type Human Deoxycytidine kinase (dCK) purified by nickel-sepharose chromatography.
Species	Human
Source	E.coli
ProteinLength	1-260 aa
Description	Deoxycytidine kinase (DCK) is required for the phosphorylation of several deoxyribonucleosides and their nucleoside analogs. Deficiency of DCK is associated with resistance to antiviral and anticancer chemotherapeutic agents. Conversely, increased deoxycytidine kinase activity is associated with increased activation of these compounds to cytotoxic nucleoside triphosphate derivatives. DCK is clinically important because of its relationship to drug resistance and sensitivity.
Tag	N-His
Molecular Mass	~31 kDa
AA Sequence	MATPPKRSCPSFSASSEGETRIKKISIEGNIAAGKSTFVNILKQLCEDWEVVPEPVARW CNVQSTQDEFEELTMSQKNGGNVLQMMYEKPERWSFTFQTYACLSRIRAQLASLN GKLKDAEKPVLFERSVYSDDRYIFASNLYESECMMNETEWTIYQDWDHWMNNQFGQ SLELDGIIYLQATPETCLHRIYLRGRNEEQGIPLEYLEKLHYKHESWLLHRTLKTNFDY

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	LQEVPILTLDVNEDFKDKYESLVEKVKE
Bio-activity	6 IU/mg protein; kcat= 0.05/sec with dC as substrate; One unit of WT human dCK converts 1.0 μ mole of dC and ATP to dCMP and ADP per minute at pH 7.5 at 37 centigrade, as measured by a coupled enzyme system with 200 μ M dC and 1 mM ATP.
Purity	> 99% (SDS-PAGE)
Storage	At -80 centigrade.
Storage Buffer	25 mM Hepes pH7.5, 200 mM NaCitrate, 10% glycerol, 5 mM DTT, 1 mM EDTA
Concentration	4.6 mg/mL
Shipping	Dry ice.
Reference	1. Sabini E, Ort S, Monnerjahn C, Konrad M, Lavie A. Structure of human dCK suggests strategies to improve anticancer and antiviral therapy. Nat Struct Biol. 2003 Jul;10(7):513-9. Sabini E, Hazra S, Konrad M, Lavie A. 2. Nonenantioselectivity property of human deoxycytidine kinase explained by structures of the enzyme in complex with L- and D-nucleosides. J Med Chem. 2007 Jun 28;50(13):3004-14. Epub 2007 May 27.

GENE INFORMATION

Gene Name	DCK deoxycytidine kinase [Homo sapiens (human)]
Official Symbol	DCK
Synonyms	DCK; deoxycytidine kinase; deoxycytidine kinase; deoxyadenosine kinase;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

deoxyguanosine kinase; deoxynucleoside kinase; EC 2.7.1.113; EC 2.7.1.74; EC 2.7.1.76

Gene ID 1633


mRNA Refseq NM_000788

Protein Refseq NP_000779

MIM 125450

UniProt ID P27707

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA