

Recombinant Human TGFB1I1 293 Cell Lysate

Cat. No. TGFB1I1-1120HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for transforming growth factor beta 1 induced transcript 1 (TGFB1I1), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

TGFB11 transforming growth factor beta 1 induced transcript 1 [Homo sapiens]

Official Symbol

TGFB11

Synonyms

TGFB11; transforming growth factor beta 1 induced transcript 1; transforming growth factor beta-1-induced transcript 1 protein; ARA55; Hic 5; HIC 5; TSC 5; androgen receptor coactivator ARA55; hydrogen peroxide-inducible clone 5 protein; androgen receptor coactivator 55 kDa protein; androgen receptor-associated protein of 55 kDa; HIC5; HIC-5; TSC-5;

Gene ID

7041

mRNA Refseq

NM_001042454

Protein Refseq

NP_001035919

MIM

602353

UniProt ID

O43294

**Chromosome
Location**

16p11

Pathway

Androgen Receptor Signaling Pathway, organism-specific biosystem; Coregulation of Androgen receptor activity, organism-specific biosystem; Monoamine Transport,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



organism-specific biosystem;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA