

Active Recombinant Human TIGIT Homodimer Protein, His tagged

Cat. No. TIGIT-60H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	TIGIT protein dimer contains a TIGIT extracellular domain (UniProt# Q495A1) with a His tag at the C-terminus was expressed in HEK293T.
Species	Human
Source	HEK293T
ProteinLength	22-141 aa
Description	Human TIGIT (T-cell immunoreceptor with Ig and ITIM domains) is also known as VSIG9 (V-set and immunoglobulin domain-containing protein 9) and VSTM3 (V-set and transmembrane domain-containing protein 3). TIGIT is a type I membrane protein containing an immunoglobulin variable (Ig-V) domain, a transmembrane domain and cytoplasmic domain. TIGIT is an immune receptor present on peripheral memory and regulatory CD4+ T cells and natural killer (NK) cells. TIGIT binds to CD155 (the poliovirus receptor, PVR) with high affinity and binds to CD112 (PVRL2) with lower affinity. Nectin-4 is also a ligand for TIGIT.
Molecular Mass	42 kDa
Purity	Greater than 90% dimer form as determined by SDS-PAGE under non-reducing condition
Application	Verified Applications: ELISA for TIGIT-specific antibody and Nectin-4 ligand protein

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

binding assays. Suggested Applications: ELISA for TIGIT-specific antibody, Nectin-4 ligand protein, and CD155 receptor protein binding assays. SPR & BLI for TIGIT-specific antibody, Nectin-4 ligand protein, and CD155 receptor protein binding assays. Animal immunization, RUO.

Storage Store at -80 centigrade.

Storage Buffer 0.22µm filtered PBS, pH 7.4

Shipping Frozen Dry Ice

GENE INFORMATION

Gene Name TIGIT T cell immunoreceptor with Ig and ITIM domains [Homo sapiens (human)]

Official Symbol TIGIT

Synonyms TIGIT; T cell immunoreceptor with Ig and ITIM domains; V set and immunoglobulin domain containing 9, V set and transmembrane domain containing 3, VSIG9, VSTM3; T-cell immunoreceptor with Ig and ITIM domains; DKFZp667A205; FLJ39873; V-set and transmembrane domain containing 3; V-set and immunoglobulin domain containing 9; Washington University cell adhesion molecule; V-set and transmembrane domain-containing protein 3; V-set and immunoglobulin domain-containing protein 9; VSIG9; VSTM3; WUCAM;

Gene ID 201633

mRNA Refseq NM_173799

Protein Refseq NP_776160

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM 612859

UniProt ID Q495A1

Bioactivity-Antibody Binding




Immobilized TIGIT-His dimer protein at 2 µg/mL (100 µL/well) can bind anti-human TIGIT monoclonal antibody with half maximal effective concentration (EC₅₀) range of 4.1-16.6 ng/mL (QC tested).

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE and WB



MW: Molecular Weight marker reduced condition NR: TIGIT dimer under non-reducing condition

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA