

Recombinant Human CXCL5 protein, mFc-tagged

Cat. No. CXCL5-3783H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CXCL5 protein(P42830)(Ala37-Asn114), fused with C-terminal mFc tag, was expressed in HEK293.
Species	Human
Source	HEK293
Form	Lyophilized from 0.22 µm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.
Molecular Mass	35.0 kDa
Endotoxin	<0.1 EU/µg protein, LAL method
Purity	>90 %, SDS-PAGE
Storage	It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. Until expiry date, -20°C to -80°C as lyophilized proteins. 3 months, -20°C to -80°C under sterile conditions after reconstitution.
Reconstitution	Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

GENE INFORMATION


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	CXCL5 chemokine (C-X-C motif) ligand 5 [Homo sapiens]
Official Symbol	CXCL5
Synonyms	CXCL5; chemokine (C-X-C motif) ligand 5; SCYB5, small inducible cytokine subfamily B (Cys X Cys), member 5 (epithelial derived neutrophil activating peptide 78); C-X-C motif chemokine 5; ENA 78; ENA-78(1-78); small inducible cytokine B5; small-inducible cytokine B5; neutrophil-activating protein 78; neutrophil-activating peptide ENA-78; epithelial-derived neutrophil activating protein 78; epithelial-derived neutrophil-activating peptide 78; epithelial-derived neutrophil-activating protein 78; small inducible cytokine subfamily B (Cys-X-Cys), member 5 (epithelial-derived neutrophil-activating peptide 78); SCYB5; ENA-78;
Gene ID	6374
mRNA Refseq	NM_002994
Protein Refseq	NP_002985
MIM	600324
UniProt ID	P42830

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA