

Active Recombinant Guinea pig IL2 protein, His-tagged

Cat. No. IL2-4743G **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Guinea pig IL2 protein(O88210)(21~152 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Guinea pig
Source	E.coli
ProteinLength	21-152 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	IL-2 (Interleukin-2) is a cytokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2 is a type of signaling molecule in the immune system, that is required for both T-cell and B-cell proliferation and other activities crucial to regulation of the immune response. Therefore, in order to detect the bioactivity of recombinant guinea pig IL-2, spleen single suspensions were prepared, activated with conA (final concentration 3 ug/ml). Cells were collected after 72h and washed with hanks. Then mouse splenic lymphocytes were seeded into triplicate wells of 96-well plates at a density of 10,000 cells/well with or without the addition of various concentrations of recombinant guinea pig IL-2. After incubated for 96h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). 10 µof CCK-8 solution was added to each well of the plate, the absorbance at 450 nm was measured using a microplate reader after incubating

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the plate for 1-4 hours at 37°C. Proliferation of Splenic lymphocytes cells after incubation with IL-2 for 96h observed by inverted microscope was shown. The dose-effect curve of recombinant guinea pig IL-2 was shown. It was obvious that recombinant guinea pig IL-2 significantly promoted cell proliferation of Splenic lymphocytes cells. The EC50 for this effect is typically 0.195- 0.323 ug/ml.

Molecular Mass	18.8kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	IL2 interleukin 2 [<i>Cavia porcellus</i>]
Official Symbol	IL2
Synonyms	IL2; interleukin 2; interleukin-2;
Gene ID	100379540
mRNA Refseq	NM_001172837
Protein Refseq	NP_001166308

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA