

Active Recombinant Zebrafish TNF protein, His-tagged

Cat. No. TNF-5513Z **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Zebrafish TNF protein(Q08CQ3)(88~234 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Zebrafish
Source	E.coli
ProteinLength	88-234 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Tumor necrosis factor (TNF, tumor necrosis factor alpha, TNFα, cachexin, orcachectin) is a cell signaling protein (cytokine) involved in systemic inflammation and is one of the cytokines that make up the acute phase reaction. Theprimaryrole of TNF is in the regulation of immune cells. TNF, being an endogenouspyrogen, is able to induce fever, apoptotic cell death, cachexia, inflammation and to inhibit tumorigenes is and viral replication and respond to sepsis viaIL1&IL6producing cells. To test the effect of TNFα on cell apoptosis, A549 cells were seeded into 96-well plates at a density of 4,000 cells/well with 5%serum standard DMEM including various concentrations of recombinant rabbit TNFα. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10μof CCK-8solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

incubating the plate for 1h at 37°C. Proliferation of A549 cells after incubation with TNFα for 72h observed by inverted microscope was shown. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant rabbit TNFα for 72h. It was obvious that TNFα significantly inhibit cell viability of A549 cells. The ED50 is 3.02 µg/mL.

Molecular Mass 20.1kDa

Endotoxin <1.0EU per 1µg (determined by the LAL method).


Purity > 90% as determined by SDS-PAGE.

Storage Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

Activity

SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA