

Active Recombinant Rabbit TNF protein, His-tagged

Cat. No. TNF-4216R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Rabbit TNF protein(P04924)(80~235 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Rabbit
Source	E.coli
ProteinLength	80-235 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Tumor necrosis factor (TNF, tumor necrosis factor alpha, TNFα, cachexin, orcachectin) is a cell signaling protein (cytokine) involved in systemic inflammation and is one of the cytokines that make up the acute phase reaction. The primaryrole of TNF is in the regulation of immune cells. TNF, being an endogenouspyrogen, is able to induce fever, apoptotic cell death, cachexia, inflammation and to inhibit tumorigenes is and viral replication and respond to sepsis vialL1&IL6 producing cells. To test the effect of TNFα on cell apoptosis, A549 cells were seeded into 96-well plates at a density of 4,000 cells/well with 5% serum standard DMEM including various concentrations of recombinant pig TNFα. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10μof CCK-8 solutionwasadded to each well of the plate, then the absorbance at 450nmwas measuredusing a microplate reader after

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

incubating the plate for 1h at 37°C. Proliferation of A549 cells after incubation with TNFα for 72h observed by inverted microscopewas shown. Cell viability was assessed by CCK-8 (Cell CountingKit-8)assay after incubation with recombinant pig TNFα for 72h. It was obvious that TNFα significantly inhibit cell viability of A549 cells. The ED50 is 10.1 µg/mL.

Molecular Mass	21.0kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 97% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	TNF tumor necrosis factor [<i>Oryctolagus cuniculus</i>]
Official Symbol	TNF
Synonyms	TNF; tumor necrosis factor; TNF-a; TNF-alpha; cachectin; tumor necrosis factor alpha; tumor necrosis factor (TNF superfamily, member 2); tumor necrosis factor ligand superfamily member 2; TNFA; TNFSF2;
Gene ID	100009088
mRNA Refseq	NM_001082263

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Protein Refseq

NP_001075732

Activity



SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA