

Active Recombinant Human LH protein, His&hFc-tagged

Cat. No. LH-2690H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human LH protein(P01215; P01229)(25~116 aa+GGGGS*3+21~141 aa), fused with N-terminal His tag and C-terminal hFc tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	25-116 aa+GGGGS*3+21-141 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Luteinizing Hormone (LH) is a 42 kDa heterodimer belonging to the glycoprotein hormone family. It is composed of noncovalently linked glycosylated alpha and beta chains. The alpha subunit (CG alpha) is also a component of Follicle-Stimulating Hormone (FSH), Thyroid-Stimulating Hormone, and Chorionic Gonadotropin. The unique beta subunit confers the protein's specific biological action and is responsible for the interaction with its receptor. LH is produced and secreted by the anterior pituitary gland. Its secretion is controlled by Gonadotropin-Releasing Hormone from the hypothalamus; however, LH secretion can also be stimulated by estradiol. A functional binding ELISA assay was conducted to detect the interaction of recombinant human CGa/LHb and recombinant human Dopamine Receptor D1 (DRD1). Briefly, biotin-linked CGa/LHb were diluted serially in PBS, with 0.01% BSA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

(pH 7.4). Duplicate samples of 100 μ l were then transferred to DRD1-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST 3 times and incubation with Streptavidin-HRP for 30min, then wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 μ stop solution to the wells and read at 450 nm immediately. The binding activity of CGa/LHb and DRD1 was shown, the EC50 for this effect is 0.34 μ g/mL.

Molecular Mass 55.3kDa


Endotoxin <1.0EU per 1 μ g (determined by the LAL method).

Purity > 90% as determined by SDS-PAGE.


Storage Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

Activity



SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA