

Active Recombinant Human PLAT protein, His-tagged

Cat. No. PLAT-3076H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human PLAT protein(P00750)(36~562 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	36-562 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Plasminogen activators are serine proteases that catalyze the activation of plasmin via proteolytic cleavage of its zymogen form plasminogen. Plasmin is an important factor in fibrinolysis, the breakdown of fibrin polymers formed during blood clotting. There are two main plasminogen activators: urokinase (uPA) and tissue plasminogen activator (tPA). Tissue plasminogen activators (tPA) are used to treat medical conditions related to blood clotting including embolic or thrombotic stroke, myocardial infarction, and pulmonary embolism. Besides, Plasminogen Activator Inhibitor 1 (PAI1) has been identified as an interactor of tPA, thus a binding ELISA assay was conducted to detect the interaction of recombinant human tPA and recombinant human PAI1. Briefly, tPA were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100ul were then transferred to PAI1-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

anti-tPA pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50μ stop solution to the wells and read at 450nm immediately. The binding activity of tPA and PAI1 was shown, and this effect was in a dose dependent manner.

Molecular Mass	60.7kDa
Endotoxin	<1.0EU per 1μg (determined by the LAL method).
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	PLAT plasminogen activator, tissue [Homo sapiens]
Official Symbol	PLAT
Synonyms	PLAT; plasminogen activator, tissue; tissue-type plasminogen activator; alteplase; reteplase; t-plasminogen activator; plasminogen activator, tissue type; tissue plasminogen activator (t-PA); TPA; T-PA; DKFZp686l03148;
Gene ID	5327
mRNA Refseq	NM_000930

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_000921


MIM 173370

UniProt ID P00750

Activity



SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA