

Recombinant Mouse Sphingosine Kinase 2, His-tagged

Cat. No. Sphk2-526M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Sphingosine kinase 2 (full length) was expressed in <i>aBaculovirus infected Sf9cell</i> expression system with N-terminal His tag. MW=68 kDa.
Species	Mouse
Source	Sf9 Cells
Description	Sphingosine kinase (SphK) is a conserved lipid kinase that catalyzes formation Sphingosine-1-phosphate (S1P) from the precursor sphingolipid sphingosine. Sphingolipid metabolites, such as ceramide, sphingosine and sphingosine-1-phosphate, are lipid second messengers involved in diverse cellular processes. There are two forms of SphK, SphK1 and SphK2. SPHK1 is found in the cytosol of eukaryotic cells, and migrates to the plasma membrane upon activation. SphK2 is localized to the nucleus.
Formulated In	25 mM Tris-HCl, pH 8.0, 100 mM NaCl, 0.05% Tween-20, 50% glycerol, and 3 mM DTT.
Application	Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.
Specific Activity	2 U/ug. Sphingosine kinase activity was determined using D-erythrosphingosine and ATP as substrates. A unit of sphingosine kinase activity is defined as the amount of enzyme required to produce 1 pmol of S1P/min. Assay buffer: 50 mM HEPES, pH 7.4, 150 mM NaCl, 5 mM MgCl ₂ , 1 mM DTT, 3 uM Na-orthovanadate, 0.5 mM ATP, 4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	uM D-erythrosphingosine, and 5 ug/ml mouse sphingosine kinase 2.
Purity	>70% by SDS-PAGE.
Stability	> 6 months at –80°C.
Pathways	Calcium signaling pathway; Fc gamma R-mediated phagocytosis; Metabolic pathways; Sphingolipid metabolism; VEGF signaling pathway
Full Length	Full L.
GENE INFORMATION	
Gene Name	Sphk2 sphingosine kinase 2 [Mus musculus]
Synonyms	sphingosine kinase 2; C76851; MGC102297; Sphk2; OTTMUSP00000026912; OTTMUSP00000026913; OTTMUSP00000026914
Gene ID	56632
mRNA Refseq	NM_020011
Protein Refseq	NP_064395
UniProt ID	Q9JIA7
Chromosome Location	7 B4
Function	ATP binding; D-erythro-sphingosine kinase activity; diacylglycerol kinase activity; kinase activity; nucleotide binding; protein binding; sphinganine kinase activity; transferase activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA