

Recombinant Human SNTA1 293 Cell Lysate

Cat. No. SNTA1-1608HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for syntrophin, alpha 1 (dystrophin-associated protein A1, 59kDa, acidic component) (SNTA1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name SNTA1 syntrophin, alpha 1 (dystrophin-associated protein A1, 59kDa, acidic component) [Homo sapiens]

Official Symbol SNTA1

Synonyms SNTA1; syntrophin, alpha 1 (dystrophin-associated protein A1, 59kDa, acidic component); SNT1, syntrophin, alpha 1 (dystrophin associated protein A1, 59kD, acidic component); alpha-1-syntrophin; pro TGF alpha cytoplasmic domain interacting protein 1; TACIP1; syntrophin-1; acidic alpha 1 syntrophin; pro-TGF-alpha cytoplasmic domain-interacting protein 1; 59 kDa dystrophin-associated protein A1 acidic component 1; SNT1; LQT12; dJ1187J4.5;

Gene ID 6640

mRNA Refseq NM_003098

Protein Refseq NP_003089

MIM 601017

UniProt ID Q13424

Chromosome Location 20q11.2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Signaling mediated by p38-gamma and p38-delta, organism-specific biosystem;
----------------	---

Function	PDZ domain binding; actin binding; calmodulin binding; protein binding;
-----------------	---

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA