

Human DLAT Knockdown Cell Lysate

Cat. No. DLAT-137HKCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	WB-validated DLAT Knockdown HeLa Cell Lysate
Species	Human
Source	HeLa
Description	This gene encodes component E2 of the multi-enzyme pyruvate dehydrogenase complex (PDC). PDC resides in the inner mitochondrial membrane and catalyzes the conversion of pyruvate to acetyl coenzyme A. The protein product of this gene, dihydrolipoamide acetyltransferase, accepts acetyl groups formed by the oxidative decarboxylation of pyruvate and transfers them to coenzyme A. Dihydrolipoamide acetyltransferase is the antigen for antimitochondrial antibodies. These autoantibodies are present in nearly 95% of patients with the autoimmune liver disease primary biliary cirrhosis (PBC). In PBC, activated T lymphocytes attack and destroy epithelial cells in the bile duct where this protein is abnormally distributed and overexpressed. PBC eventually leads to cirrhosis and liver failure. Mutations in this gene are also a cause of pyruvate dehydrogenase E2 deficiency which causes primary lactic acidosis in infancy and early childhood.
Form	Cell-Tissue Lysis buffer
Molecular Mass	69 kDa
Notes	Instruction of use: This knockdown cell lysate should be paired with wild-type HeLa cell lysate for use. For Western blotting, we recommend running wild-type and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	knockdown lysates on the same gel, and loading each well with equal volume and equal amount of total proteins.
Storage	Store at -20 centigrade for two years.
Concentration	Lot-specific
Shipping	Blue Ice
Components	1 vial of 100 µg WT HeLa cell lysate 1 vial of 100 µg DLAT KD HeLa cell lysate
Protein Families	Druggable Genome
Protein Pathways	Citrate cycle (TCA cycle), Glycolysis / Gluconeogenesis, Metabolic pathways, Pyruvate metabolism
Lysate QC	RT-qPCR; Western Blotting (WB)

GENE INFORMATION

Gene Name	DLAT dihydrolipoamide S-acetyltransferase [Homo sapiens (human)]
Official Symbol	DLAT
Synonyms	DLAT; dihydrolipoamide S-acetyltransferase; DLTA; dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex, mitochondrial; E2 component of pyruvate dehydrogenase complex; PDC E2; PBC; M2 antigen complex 70 kDa subunit; pyruvate dehydrogenase complex component E2; 70 kDa mitochondrial autoantigen of primary biliary cirrhosis; dihydrolipoamide acetyltransferase component of pyruvate dehydrogenase complex; dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

complex mitochondrial; PDCE2; PDC-E2;

Gene ID 1737**mRNA Refseq** NM_001931**Protein Refseq** NP_001922**MIM** 608770**UniProt ID** P10515 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA