

Recombinant Human SLC43A2 293 Cell Lysate

Cat. No. SLC43A2-1713HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for solute carrier family 43, member 2 (SLC43A2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [SLC43A2 solute carrier family 43, member 2 \[Homo sapiens \]](#)

Official Symbol SLC43A2

Synonyms SLC43A2; solute carrier family 43, member 2; large neutral amino acids transporter small subunit 4; MGC34680; amino acid transporter; L-type amino acid transporter 4; LAT4; FLJ23848;

Gene ID [124935](#)

mRNA Refseq [NM_152346](#)

Protein Refseq [NP_689559](#)

MIM [610791](#)

UniProt ID [Q8N370](#)

Chromosome Location 17p13.3

Pathway Amino acid transport across the plasma membrane, organism-specific biosystem; SLC-mediated transmembrane transport, organism-specific biosystem; Transmembrane transport of small molecules, organism-specific biosystem; Transport of inorganic cations/anions and amino acids/oligopeptides, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem;

Function

L-amino acid transmembrane transporter activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA