

Active Native Bovine Lactoperoxidase

Cat. No. Lactoperoxidase-18B **Lot. No.** (See product label)

SPECIFICATION

Product Overview	LPO is a glycoprotein with a single hemin prosthetic group per molecule. It may consist of two isozymes.
Species	Bovine
Source	Milk
Description	Lactoperoxidase (LPO) is a glycoprotein with a hemin prosthetic group which may occur as a mixture of two isozymes. It has a molecular weight of 77,500 daltons. LPO catalyzes the hydrogen peroxide oxidation of iodide according to the following reaction: $2I^- + H_2O_2 + 2H^+ \rightarrow I_2 + 2H_2O$. Iodide reacts directly with the heme group; upon addition of H_2O_2 the complex iodates the substrate. LPO is inhibited by hydrazines. The assay procedure has been updated from that of Morrison to an ABTS®/ H_2O_2 based method with increased sensitivity and reproducibility.
Form	A lyophilized powder.
Bio-activity	≥35 units/mg dry weight ABTS
Molecular Mass	77.5 kDa
Purity	Chromatographically purified. $A_{412}/A_{280} = 0.93-0.96$
Storage	Store at -20 centigrade.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

EC	1.11.1.7
CAS	9003-99-0
Unit definition	One Unit reduces one micromole of hydrogen peroxide per minute at 25 centigrade, pH 6.0. Note: This unit definition is based upon a new assay method. The previous method yielded approximately 2.8 times higher results.
Usage	Method: The assay procedure is a modification of that described by Putter and Becker. The increase in the absorbance at 405 nm resulting from the oxidation of 2,2-azino-di(3-ethyl-benzothiazoline-6-sulphonic acid), diammonium salt, ABTS. Two moles of ABTS are oxidized for every mole of peroxide that is reduced. Reagents: 0.08 M Sodium-Potassium Phosphate Buffer pH 6.0: Prepare by adjusting the pH of 0.08 M sodium phosphate, dibasic, to 6.0 using 0.08 M potassium phosphate, monobasic. Sodium Potassium Phosphate Buffer, containing 1 mg/mL Bovine Serum Albumin: Dissolve BSA at a concentration of mg/ml in an aliquot of the above buffer. This will be used as enzyme diluent. 0.0017 M Hydrogen peroxide. Prepare by diluting 1 ml of 30% hydrogen peroxide to 100 ml with reagent grade water. Further dilute 1 ml of this solution to 50 ml with reagent grade water. Prepare fresh daily. 0.051 M ABTS: Dissolve ABTS in 0.08 M sodium phosphate buffer, pH 6.0 Enzyme: Dissolve at one mg/ml in sodium-phosphate buffer. Immediately prior to use, dilute further in phosphate buffer with BSA to obtain a rate of 0.008-0.025 DA/minute. Keep all dilutions on ice. Procedure: Set spectrophotometer at 25 centigrade and 405 nm. Pipette into each cuvette 2.65 ml sodium-phosphate buffer, 0.15 mL hydrogen peroxide, and 0.10 ml ABTS. Incubate in spectrophotometer at 25 centigrade for 3-5 minutes to achieve temperature equilibration and establish a blank rate, if any. Add 0.10 ml diluted enzyme and record the increase in A405 for 4-6 minutes. Calculate the rate from the maximum initial linear portion of the curve. The reaction is linear for approximately two minutes. Calculation: (Units/mg)=($\Delta A_{350}/\text{min} \times 3.0 \times \text{dilution}$)/(18.6$\times 2 \times \text{sample volume}$) 18.6= extinction coefficient of ABTS at 405nm 2 ABTS: 1 H₂O₂

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Extinction
Coefficient**

13.9

Inhibitors

LPO is inhibited by hydrazines. Dolman et al. (1968) report on the kinetics of cyanide inhibition.

GENE INFORMATION**Official Symbol**

Lactoperoxidase

Synonyms

Lactoperoxidase

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA