

Active Recombinant Human CCL21 protein, His-tagged

Cat. No. CCL21-10969H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human CCL21 Protein, His Tag is expressed from Baculovirus-Insect cells. It contains AA Ser 24-Pro 134 (Accession # O00585). This protein carries a polyhistidine tag at the C-terminus. Predicted N-terminus: Ser 24
Species	Human
Source	Insect Cells
ProteinLength	24-134 aa
Description	CC chemokine 21 (CCL21), also known as secondary lymphoidorgan chemokine (SLC), is a ligand for the chemokine receptor CCR7. CCL19 and CCL21 are the only ligands for CCR7, which is expressed by various subsets of immune cells. CCL19 and CCL21 function to recruit CCR7 expressing antigen presenting cells, like dendritic cells and naive T-cells, to the lymph nodes. CCL21 seems to be the critical CCR7 ligand regulating dendritic cell homeostasis and function. CCL21 has an unusual extended C-terminal tail for a chemokine. Only one gene exists for this chemokine in human beings, whereas mice have two. CCL21 Inhibit hemopoiesis and stimulates chemotaxis. Chemotactic in vitro for thymocytes and activated T-cells, but not for B-cells, macrophages, or neutrophils.
Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Bio-activity	Immobilized Human CCL21 Protein, His Tag at 5 µg/mL (100 µL/well) can bind Human CCR7 Full Length Protein, His, Flag Tag with a linear range of 0.01-0.313

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	µg/mL.
Molecular Mass	The protein has a calculated MW of 14.1 kDa. The protein migrates as 17-19 kDa when calibrated against Star Ribbon Pre-stained Protein Marker under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 0.1 EU/µg by the LAL method/rFC method.
Purity	>90% as determined by SDS-PAGE.
Storage	This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution. For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.

GENE INFORMATION

Gene Name	CCL21 chemokine (C-C motif) ligand 21 [Homo sapiens (human)]
Official Symbol	CCL21
Synonyms	CCL21; chemokine (C-C motif) ligand 21; SCYA21, small inducible cytokine subfamily A (Cys Cys), member 21; C-C motif chemokine 21; 6Ckine; beta chemokine exodus 2; CKb9; ECL; Efficient Chemoattractant for Lymphocytes; exodus 2; secondary lymphoid tissue chemokine; SLC; TCA4; exodus-2; beta chemokine exodus-2; beta-chemokine exodus-2; small-inducible cytokine A21; secondary lymphoid-tissue chemokine; small inducible cytokine subfamily A (Cys-Cys), member 21; SCYA21;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	MGC34555;
Gene ID	6366
mRNA Refseq	NM_002989
Protein Refseq	NP_002980
MIM	602737
UniProt ID	O00585

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA