

Active Recombinant Monkey HAVCR2 Protein, Fc-tagged

Cat. No. HAVCR2-11155M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Cynomolgus TIM-3, Fc Tag is expressed from human 293 cells (HEK293). It contains AA Ser 22 - Arg 201 (Accession # EHH54703.1). This protein carries a human IgG1 Fc tag at the C-terminus. Predicted N-terminus: Ser 22
Species	Monkey
Source	HEK293
ProteinLength	22- 201 aa
Description	Hepatitis A virus cellular receptor 2 is also known as HAVCR2, FLJ14428, KIM3, TIM3, TIMD3, is a member of the TIM family of immune regulating molecules with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. CD4-positive T helper lymphocytes can be divided into types 1 (Th1) and 2 (Th2) on the basis of their cytokine secretion patterns. Th1 cells and their associated cytokines are involved in cell-mediated immunity to intracellular pathogens and delayed-type hypersensitivity reactions, whereas Th2 cells are involved in the control of extracellular helminthic infections and the promotion of atopic and allergic diseases. The 2 types of cells also cross-regulate the functions of the other. HAVCR2 is a Th1-specific cell surface protein that regulates macrophage activation and enhances the severity of experimental autoimmune encephalomyelitis in mice. HAVCR2 regulates macrophage activation. Inhibits T-helper type 1 lymphocyte (Th1)-mediated auto- and alloimmune responses and promotes immunological tolerance. May be also involved in T-cell homing. Dysregulation of the HAVCR2-galectin-9 pathway could underlie chronic

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	autoimmune disease states in human, such as multiple sclerosis.
Form	Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.
Bio-activity	Immobilized Cynomolgus TIM-3, Fc Tag at 2 µg/mL (100 µL/well) can bind Anti-TIM3 Mab, Human IgG4 with a linear range of 0.5-8 ng/mL.
Molecular Mass	The protein has a calculated MW of 45.8 kDa. The protein migrates as 50-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU/µg by the LAL method/rFC method.
Purity	>85% as determined by SDS-PAGE.
Storage	This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution. For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.

GENE INFORMATION

Gene Name	HAVCR2 hepatitis A virus cellular receptor 2 [Macaca mulatta (Rhesus monkey)]
Official Symbol	HAVCR2
Synonyms	Havcr2; hepatitis A virus cellular receptor 2; Tim3; TIM-3; Timd3; hepatitis A virus

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cellular receptor 2 homolog; HAVcr-2; T-cell immunoglobulin and mucin domain containing 3; T-cell immunoglobulin and mucin domain-containing protein 3; T-cell immunoglobulin mucin receptor 3; T-cell membrane protein 3; TIMD-3

Gene ID [714891](#)

mRNA Refseq [XM_015141316](#)

Protein Refseq [XP_014996802](#)

UniProt ID [F7GB92](#)

SDS-PAGE.

