Recombinant Human CD7 Protein, His tagged

Cat. No. CD7-122H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD7 (Ala26-Pro180) with C-terminus His tag is expressed in CHO cell
Species	Human
Source	CHO
Conjugation/Label	26-180 aa
Description	CD7 is a transmembrane glycoprotein belonging to the immunoglobulin superfamily that serves as an important marker of T cell and natural killer (NK) cell lineages. It is one of the earliest surface antigens expressed during T cell development, appearing on thymocytes and persisting throughout T cell maturation. CD7 plays a costimulatory role in T cell activation and is involved in regulating immune responses. The protein participates in T cell receptor signaling, cell-cell interactions, and may contribute to immune synapse formation and lymphocyte trafficking, though its complete physiological functions continue to be investigated. Structurally, CD7 is a type I transmembrane protein of approximately 40 kDa consisting of a single extracellular immunoglobulin-like domain, a transmembrane region, and a cytoplasmic tail. The extracellular domain contains the characteristic features of the immunoglobulin superfamily, including disulfide bonds that stabilize its three-dimensional structure. The cytoplasmic domain is relatively short but contains motifs that can interact with intracellular signaling molecules and cytoskeletal components. CD7 can form homodimers on the cell surface, and this oligomerization may be important for its

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	signaling and costimulatory functions. The primary identified ligand for CD7 is K12/SECTM1 (secreted and transmembrane 1), which binds to CD7 and mediates costimulatory signals during T cell activation. Upon binding to K12/SECTM1, CD7 triggers the production of cytokines and enhances T cell responses. Additionally, CD7 may engage in homophilic interactions (CD7-CD7 binding between cells) and has been reported to interact with galectin-1, a lectin involved in immune regulation. The protein's glycosylation patterns influence its binding properties and functional activities in the immune system. In disease contexts, CD7 expression patterns are clinically significant in hematologic malignancies. CD7 is expressed on most T cell acute lymphoblastic leukemias (T-ALL) and certain peripheral T cell lymphomas, making it a valuable diagnostic marker. CD7-positive acute myeloid leukemia (AML) represents a subset with distinct clinical features and often poorer prognosis. Therapeutically, CD7 has emerged as a promising target for immunotherapy in T cell malignancies. CD7-directed chimeric antigen receptor (CAR) T cell therapies are under active development and clinical investigation for treating T-ALL and peripheral T cell lymphomas. However, targeting CD7 presents unique challenges because normal T cells also express this marker, requiring innovative strategies such as CD7 knockout in CAR-T cells to prevent fratricide (self-destruction). K12/SECTM1-based CAR T cells that exploit the natural CD7-K12 interaction are being developed to enhance specificity and efficacy while minimizing toxicity to normal T cells.
Form	Liquid, 1×PBS buffer, pH7.4, 0.22 μm filtered
Molecular Mass	The protein has a predicted molecular weight of 16 kDa. Under DTT-reducing conditions, it migrates at approximately 30-40 kDa on SDS-PAGE.
Purity	>95% based on SDS-PAGE under reducing condition
Applications	ELISA, BLI

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA


Storage	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4 centigrade for up to 2 weeks, or at -20 centigrade or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 μm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ –70 centigrade to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.
Concentration	25 μg size is bottled at 0.2 mg/mL concentration. 100 μg size is supplied at a lot-specific concentration.

GENE INFORMATION

Gene Name	CD7 CD7 molecule [Homo sapiens (human)]
Official Symbol	CD7
Synonyms	CD7; CD7 molecule; CD7 antigen (p41); T-cell antigen CD7; GP40; LEU 9; p41 protein; T cell antigen CD7; T cell leukemia antigen; Tp40; TP41; T-cell leukemia antigen; T-cell surface antigen Leu-9; LEU-9;
Gene ID	924
mRNA Refseq	NM_006137
Protein Refseq	NP_006128
MIM	186820
UniProt ID	P09564


Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



Human CD7 Protein (C-His) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The gel was stained for 1 hour with BlinkBlue. The purity of this protein appears to be greater than 95% based on reducing conditions.

ELISA



Human SECTM1 (C-Fc) protein is coated at 1 μg/mL (100 ng/well). Human CD7 Protein (C-His) can detect the SECTM1 protein in the dose dependent manner. The ED50 is about 4-20 ng/mL.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA