

Recombinant Human SLC22A8 293 Cell Lysate

Cat. No. SLC22A8-1790HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for solute carrier family 22 (organic anion transporter), member 8 (SLC22A8) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	SLC22A8 solute carrier family 22 (organic anion transporter), member 8 [Homo sapiens]
Official Symbol	SLC22A8
Synonyms	SLC22A8; solute carrier family 22 (organic anion transporter), member 8; solute carrier family 22 member 8; OAT3; hOAT3; organic anion transporter 3; MGC24086;
Gene ID	9376
mRNA Refseq	NM_001184736
Protein Refseq	NP_001171665
MIM	607581
UniProt ID	Q8TCC7
Chromosome Location	11q12.3
Pathway	Bile secretion, organism-specific biosystem; Bile secretion, conserved biosystem; Organic anion transport, organism-specific biosystem; Organic cation/anion/zwitterion transport, organism-specific biosystem; SLC-mediated transmembrane transport, organism-specific biosystem; Transmembrane transport of small molecules,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem; Transport of glucose and other sugars, bile salts and organic acids, metal ions and amine compounds, organism-specific biosystem;

Function

inorganic anion exchanger activity; ion transmembrane transporter activity; organic anion transmembrane transporter activity; protein kinase C binding; quaternary ammonium group transmembrane transporter activity; transmembrane transporter activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA