

Recombinant Human SLC1A4 293 Cell Lysate

Cat. No. SLC1A4-1798HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for solute carrier family 1 (glutamate/neutral amino acid transporter), member 4 (SLC1A4), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

SLC1A4 solute carrier family 1 (glutamate/neutral amino acid transporter), member 4 [Homo sapiens]

Official Symbol

SLC1A4

Synonyms

SLC1A4; solute carrier family 1 (glutamate/neutral amino acid transporter), member 4; neutral amino acid transporter A; alanine/serine/cysteine/threonine transporter; ASCT1; SATT; ASCT-1; solute carrier family 1 member 4; glutamate/neutral amino acid transporter; alanine/serine/cysteine/threonine transporter 1;

Gene ID

6509

mRNA Refseq

NM_001193493

Protein Refseq

NP_001180422

MIM

600229

UniProt ID

P43007

Chromosome Location

2p15-p13

Pathway

Amino acid transport across the plasma membrane, organism-specific biosystem; SLC-mediated transmembrane transport, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Transmembrane transport of small molecules, organism-specific biosystem; Transport of inorganic cations/anions and amino acids/oligopeptides, organism-specific biosystem;

Function

L-alanine transmembrane transporter activity; L-cystine transmembrane transporter activity; NOT L-cystine transmembrane transporter activity; L-glutamine transmembrane transporter activity; L-hydroxyproline transmembrane transporter activity; L-proline transmembrane transporter activity; L-serine transmembrane transporter activity; L-threonine transmembrane transporter activity; chloride channel activity; sodium:dicarboxylate symporter activity; symporter activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA