

Recombinant Human TNFRSF17 Protein, Fc tagged, PE Labeled

Cat. No. TNFRSF17-295H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human BCMA protein (NP_001183.2) (Met1-Ala54) with C-terminus Human IgG1 Fc tag is expressed in HEK293 cells and conjugated to PE.
Species	Human
Source	HEK293
ProteinLength	PE
Conjugation/Label	1-54 aa aa
Description	B-cell maturation antigen (BCMA), also known as CD269 or TNFRSF17, is a transmembrane glycoprotein that serves as a critical regulator of B cell development and function. It belongs to the tumor necrosis factor receptor (TNFR) superfamily and is predominantly expressed on plasma cells and a subset of late-stage B cells. BCMA's primary biological role is to promote the survival, differentiation, and long-term maintenance of antibody-producing plasma cells by mediating signals from specific ligands in the TNF family. Structurally, BCMA is a type I transmembrane protein consisting of an extracellular cysteine-rich domain responsible for ligand binding, a single transmembrane region, and a short cytoplasmic tail that interacts with intracellular signaling molecules. The cytoplasmic domain lacks death domains but recruits TRAF (TNF receptor-associated factor) adaptor proteins to activate downstream pathways such as NF-κB and MAPK. These signaling cascades

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

enhance plasma cell survival and immunoglobulin production, contributing to humoral immunity maintenance. Soluble BCMA, generated through proteolytic cleavage by γ -secretase, can act as a decoy receptor to regulate ligand availability in the serum. BCMA binds two main ligands: B-cell activating factor (BAFF, also known as BLyS) and a proliferation-inducing ligand (APRIL). Both ligands are produced by myeloid and stromal cells and support B cell homeostasis. Among these, APRIL binds BCMA with higher affinity and is the primary mediator of BCMA-dependent signaling in plasma cells. The BAFF/APRIL–BCMA axis thus serves as a crucial checkpoint for sustained antibody production and plasma cell survival. In disease contexts, BCMA is strongly implicated in multiple myeloma (MM) and certain B-cell lymphomas. Its selective overexpression on malignant plasma cells makes it an important diagnostic marker and a therapeutic target. BCMA-directed treatments have revolutionized therapy for multiple myeloma, including antibody-drug conjugates (e.g., belantamab mafodotin), bispecific T cell engagers (e.g., teclistamab, elranatamab), and CAR-T cell therapies (e.g., idecabtagene vicleucel, ciltacabtagene autoleucel). These agents exploit BCMA's restricted expression pattern to deliver targeted cytotoxicity, leading to durable responses in refractory disease. Consequently, BCMA has emerged as a prototypical target in the development of next-generation immunotherapies for hematologic malignancies.

Form	Liquid, 1×PBS buffer, pH7.4, 0.09% NaN ₃ with a carrier protein
Molecular Mass	The protein has a predicted molecular weight of 33kDa. Under DTT-reducing conditions, it migrates at approximately 35-45 kDa on SDS-PAGE prior to conjugation.
Applications	Flow Cytometry
Storage	Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2–8 centigrade for up to six months.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Concentration 25 µg size is bottled at 0.1 mg/mL concentration. 100 µg size is bottled at lot specific concentration.

GENE INFORMATION

Gene Name TNFRSF17 tumor necrosis factor receptor superfamily, member 17 [Homo sapiens (human)]

Official Symbol TNFRSF17

Synonyms TNFRSF17; tumor necrosis factor receptor superfamily, member 17; BCMA; tumor necrosis factor receptor superfamily member 17; BCM; CD269; B-cell maturation factor; B cell maturation antigen; B-cell maturation protein;

Gene ID 608

mRNA Refseq NM_001192


Protein Refseq NP_001183

MIM 109545


UniProt ID Q02223

FC

**A: BCMA CAR-transfected
Stained with PE-BCMA-Fc**



**B: Mock-transfected
Stained with PE-BCMA-Fc**



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

CHO cells transfected with either BCMA CAR or Mock plasmid were stained with PE conjugated BCMA (C-Fc) protein at 4 µg/test

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA