Recombinant Human SKP1 293 Cell Lysate

Cat. No. SKP1-1813HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	HEK293
Description	Antigen standard for S-phase kinase-associated protein 1 (SKP1), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	SKP1 S-phase kinase-associated protein 1 [Homo sapiens]
Official Symbol	SKP1
Synonyms	SKP1; S-phase kinase-associated protein 1; S phase kinase associated protein 1A (p19A) , SKP1A; EMC19; MGC34403; OCP II; OCP2; p19A; TCEB1L; SIII; OCP-2; p19skp1; organ of Corti protein 2; organ of Corti protein II; cyclin A/CDK2-associated p19; transcription elongation factor B; cyclin A/CDK2-associated protein p19; cyclin-A/CDK2-associated protein p19; RNA polymerase II elongation factor-like protein OCP2; SKP1A; OCP-II;
Gene ID	6500
mRNA Refseq	NM_006930
Protein Refseq	NP_008861
MIM	601434
UniProt ID	P63208
Chromosome Location	5q31
Pathway	APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Antigen processing: Ubiquitination and Proteasome degradation, organism-specific biosystem; Canonical Wnt signaling pathway, organism-specific biosystem; Cell Cycle, organism-specific biosystem;
Function	protein binding; ubiquitin-protein ligase activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA