

Recombinant Human SFRS7 293 Cell Lysate

Cat. No. SRSF7-1902HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for splicing factor, arginine/serine-rich 7, 35kDa (SFRS7) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [SRSF7 serine/arginine-rich splicing factor 7 \[Homo sapiens \]](#)

Official Symbol SRSF7

Synonyms 9G8; AAG3; SFRS7; serine/arginine-rich splicing factor 7; splicing factor 9G8; SR splicing factor 7; aging-associated protein 3; splicing factor, arginine/serine-rich 7, 35kDa

Gene ID [6432](#)

mRNA Refseq [NM_001031684](#)

Protein Refseq [NP_001026854](#)

MIM [600572](#)

UniProt ID [Q16629](#)

Chromosome Location 2p22.1

Pathway Cleavage of Growing Transcript in the Termination Region, organism-specific biosystem; RNA Polymerase II Transcription Termination, organism-specific biosystem; Transport of Mature mRNA derived from an Intron-Containing Transcript, organism-specific biosystem

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

nucleotide binding; poly(A) RNA binding; protein binding

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA