Recombinant Human SFPQ 293 Cell Lysate

Cat. No. SFPQ-1911HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	HEK293
Description	Antigen standard for splicing factor proline/glutamine-rich (polypyrimidine tract binding protein associated) (SFPQ) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	SFPQ splicing factor proline/glutamine-rich [Homo sapiens]
Official Symbol	SFPQ
Synonyms	SFPQ; splicing factor proline/glutamine-rich; splicing factor proline/glutamine rich (polypyrimidine tract binding protein associated); splicing factor, proline- and glutamine-rich; polypyrimidine tract binding protein associated; PSF; hPOMp100; 100 kDa DNA-pairing protein; PTB-associated splicing factor; PTB-associated-splicing factor; DNA-binding p52/p100 complex, 100 kDa subunit; polypyrimidine tract-binding protein-associated splicing factor; polypyrimidine tract-binding protein-associated-splicing factor; splicing factor proline/glutamine rich (polypyrimidine tract-binding protein-associated); POMP100; DKFZp547C228; DKFZp667K0521; DKFZp686K1282;
Gene ID	6421
mRNA Refseq	NM_005066
Protein Refseq	NP_005057
MIM	605199
UniProt ID	P23246
Chromosome	1p34.3

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	mRNA processing, organism-specific biosystem;
Function	DNA binding; RNA binding; nucleotide binding; protein binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA