

Recombinant Human SAR1B 293 Cell Lysate

Cat. No. SAR1B-2063HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for SAR1 homolog B (<i>S. cerevisiae</i>) (SAR1B), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

SAR1B SAR1 homolog B (*S. cerevisiae*) [*Homo sapiens*]

Official Symbol

SAR1B

Synonyms

AR1B; SAR1 homolog B (*S. cerevisiae*); ANDD; CMRD; GTBPB; SARA2; GTP-binding protein SAR1b; 2310075M17Rik; SAR1a gene homolog 2; GTP-binding protein B; GTP-binding protein Sara; SAR1B; EC 3.6.5

Gene ID

51128

mRNA Refseq

NM_001033503

Protein Refseq

NP_001028675

MIM

607690

UniProt ID

Q9Y6B6

Chromosome Location

5q31.1

Pathway

Adaptive Immune System; COPII complex; Lipid digestion; Post-translational protein modification; Protein processing in endoplasmic reticulum; Transport to the Golgi and subsequent modification; MHC class II antigen presentation

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

GTP binding; GTPase activity; metal ion binding

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA