

Recombinant Human RRP8 293 Cell Lysate

Cat. No. RRP8-2140HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for ribosomal RNA processing 8, methyltransferase, homolog (yeast) (RRP8) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	RRP8 ribosomal RNA processing 8, methyltransferase, homolog (yeast) [Homo sapiens]
Official Symbol	RRP8
Synonyms	RRP8; ribosomal RNA processing 8, methyltransferase, homolog (yeast); KIAA0409; ribosomal RNA-processing protein 8; RRP8 methyltransferase homolog (S. cerevisiae); nucleomethylin; cerebral protein 1; RRP8 methyltransferase homolog;
Gene ID	23378
mRNA Refseq	NM_015324
Protein Refseq	NP_056139
UniProt ID	O43159
Chromosome Location	11p15.4
Function	S-adenosylmethionine-dependent methyltransferase activity; methylated histone residue binding; protein binding; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA