# Recombinant Human RPSA 293 Cell Lysate

**Cat. No.** RPSA-2155HCL **Lot. No.** (See product label)

## SPECIFICATION

| | |
|---|---|
| **Product Overview** | |
| **Species** | Human |
| **Source** | HEK293 |
| **Description** | Antigen standard for ribosomal protein SA (RPSA), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection. |
| **Components** | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| **Size** | 0.1 mg |
| **Storage Instruction** | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment. |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| **Applications** | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane. |

## GENE INFORMATION

| | |
|---|---|
| **Gene Name** | RPSA ribosomal protein SA [ Homo sapiens ] |
| **Official Symbol** | RPSA |
| **Synonyms** | SA; LBP; LRP; p40; 67LR; ICAS; lamR; 37LRP; LAMBR; LAMR1; LRP/LR; LBP/p40; NEM/1CHD4; 40S ribosomal protein SA; 37 kDa laminin receptor; 67 kDa laminin receptor; 37/67 kDa laminin receptor; laminin-binding protein precursor p40; colon carcinoma laminin-binding protein; laminin receptor 1 (67kD, ribosomal protein SA); multidrug resistance-associated protein MGr1-Ag |
| **Gene ID** | 3921 |
| **mRNA Refseq** | NM_002295 |
| **Protein Refseq** | NP_002286 |
| **MIM** | 150370 |
| **UniProt ID** | P08865 |
| **Chromosome Location** | 3p22.2 |
| **Pathway** | Activation of the mRNA upon binding of the cap-binding complex and eIFs, and subsequent binding to 43S, organism-specific biosystem; Eukaryotic Translation |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| | Elongation, organism-specific biosystem |
| **Function** | laminin binding; protein binding; structural constituent of ribosome |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA