

Recombinant Human RPS27L 293 Cell Lysate

Cat. No. RPS27L-2164HCL Lot. No. (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for ribosomal protein S27-like (RPS27L) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [RPS27L ribosomal protein S27-like \[Homo sapiens \]](#)

Official Symbol RPS27L

Synonyms RPS27L; ribosomal protein S27-like; 40S ribosomal protein S27-like;

Gene ID [51065](#)

mRNA Refseq [NM_015920](#)

Protein Refseq [NP_057004](#)

MIM [612055](#)

UniProt ID [Q71UM5](#)

Chromosome Location 15q21.3

Pathway Direct p53 effectors, organism-specific biosystem; Ribosome, organism-specific biosystem; Ribosome, conserved biosystem; Ribosome, eukaryotes, organism-specific biosystem;

Function cysteine-type endopeptidase activator activity involved in apoptotic process; metal ion binding; protein binding; structural constituent of ribosome; translation activator

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA