

Recombinant Human RPL30 293 Cell Lysate

Cat. No. RPL30-2207HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for ribosomal protein L30 (RPL30) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [RPL30 ribosomal protein L30 \[Homo sapiens \]](#)

Official Symbol RPL30

Synonyms RPL30; ribosomal protein L30; 60S ribosomal protein L30; L30;

Gene ID [6156](#)

mRNA Refseq [NM_000989](#)

Protein Refseq [NP_000980](#)

MIM [180467](#)

UniProt ID [P62888](#)

Chromosome Location 8q22

Pathway Cap-dependent Translation Initiation, organism-specific biosystem; Cytoplasmic Ribosomal Proteins, organism-specific biosystem; Disease, organism-specific biosystem; Eukaryotic Translation Elongation, organism-specific biosystem; Eukaryotic Translation Initiation, organism-specific biosystem; Eukaryotic Translation Termination, organism-specific biosystem; Formation of a pool of free 40S subunits, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

RNA binding; structural constituent of ribosome; structural constituent of ribosome;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA