

Recombinant Human RFC2 293 Cell Lysate

Cat. No. RFC2-2412HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for replication factor C (activator 1) 2, 40kDa (RFC2), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

RFC2 replication factor C (activator 1) 2, 40kDa [Homo sapiens]

Official Symbol

RFC2

Synonyms

RFC2; replication factor C (activator 1) 2, 40kDa; replication factor C (activator 1) 2 (40kD); replication factor C subunit 2; A1; activator 1; RFC40; A1 40 kDa subunit; RF-C 40 kDa subunit; activator 1 subunit 2; replication factor C 2; activator 1 40 kDa subunit; MGC3665;

Gene ID

5982

mRNA Refseq

NM_002914

Protein Refseq

NP_002905

MIM

600404

UniProt ID

P35250

Chromosome Location

7q11.23

Pathway

Activation of ATR in response to replication stress, organism-specific biosystem; BRCA1-associated genome surveillance complex (BASC), organism-specific biosystem; Cell Cycle, organism-specific biosystem; Cell Cycle Checkpoints,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



organism-specific biosystem; Cell Cycle, Mitotic, organism-specific biosystem;
Chromosome Maintenance, organism-specific biosystem; DNA Repair, organism-
specific biosystem;

Function

ATP binding; DNA clamp loader activity; nucleoside-triphosphatase activity;
nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA