

Recombinant Human RAG2 293 Cell Lysate

Cat. No. RAG2-2547HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for recombination activating gene 2 (RAG2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

RAG2 recombination activating gene 2 [Homo sapiens]

Official Symbol

RAG2

Synonyms

RAG2; recombination activating gene 2; V(D)J recombination-activating protein 2; RAG-2;

Gene ID

5897

mRNA Refseq

NM_000536

Protein Refseq

NP_000527

MIM

179616

UniProt ID

P55895

Chromosome Location

11p13

Pathway

C-MYB transcription factor network, organism-specific biosystem; Primary immunodeficiency, organism-specific biosystem; Primary immunodeficiency, conserved biosystem;

Function

DNA binding; chromatin binding; endonuclease activity; metal ion binding; methylated

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

histone residue binding; phosphatidylinositol binding; phosphatidylinositol-3,4,5-trisphosphate binding; phosphatidylinositol-3,4-bisphosphate binding; phosphatidylinositol-3,5-bisphosphate binding; phosphatidylinositol-4,5-bisphosphate binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA