

Recombinant Human RAD23B 293 Cell Lysate

Cat. No. RAD23B-2558HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for RAD23 homolog B (<i>S. cerevisiae</i>) (RAD23B) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[RAD23B](#) [RAD23 homolog B \(S. cerevisiae\)](#) [[Homo sapiens](#)]

Official Symbol

[RAD23B](#)

Synonyms

[RAD23B](#); [RAD23 homolog B \(S. cerevisiae\)](#); [RAD23 \(S. cerevisiae\) homolog B](#); UV excision repair protein [RAD23 homolog B](#); [HHR23B](#); [HR23B](#); [P58](#); XP C repair complementing complex 58 kDa; XP C repair complementing protein; [RAD23](#), yeast homolog of, B; XP-C repair complementing protein; XP-C repair complementing complex 58 kDa; XP-C repair-complementing complex 58 kDa protein;

Gene ID

[5887](#)

mRNA Refseq

[NM_001244713](#)

Protein Refseq

[NP_001231642](#)

MIM

[600062](#)

UniProt ID

[P54727](#)

Chromosome Location

9q31.2

Pathway

DNA Damage Recognition in GG-NER, organism-specific biosystem; DNA Repair, organism-specific biosystem; Dual incision reaction in GG-NER, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Formation of incision complex in GG-NER, organism-specific biosystem; Global Genomic NER (GG-NER), organism-specific biosystem; Nucleotide Excision Repair, organism-specific biosystem; Nucleotide excision repair, organism-specific biosystem;

Function

damaged DNA binding; polyubiquitin binding; protein binding; single-stranded DNA binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA