

Recombinant Human PTPN7 293 Cell Lysate

Cat. No. PTPN7-2681HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for protein tyrosine phosphatase, non-receptor type 7 (PTPN7), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[PTPN7 protein tyrosine phosphatase, non-receptor type 7 \[Homo sapiens \]](#)

Official Symbol

PTPN7

Synonyms

PTPN7; protein tyrosine phosphatase, non-receptor type 7; tyrosine-protein phosphatase non-receptor type 7; HEPTP; LC PTP; dual specificity phosphatase 1; protein-tyrosine phosphatase LC-PTP; hematopoietic protein-tyrosine phosphatase; protein-tyrosine phosphatase, nonreceptor-type, stress induced; LPTP; PTPNI; BPTP-4; LC-PTP; FLJ18703;

Gene ID

[5778](#)

mRNA Refseq

[NM_080588](#)

Protein Refseq

[NP_542155](#)

MIM

[176889](#)

UniProt ID

[P35236](#)

**Chromosome
Location**

1q32.1

Pathway

Downstream signaling in naive CD8+ T cells, organism-specific biosystem; MAPK signaling pathway, organism-specific biosystem; MAPK signaling pathway, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem; MAPK signaling pathway, conserved biosystem; Ras signaling in the CD4+ TCR pathway, organism-specific biosystem;

Function

hydrolase activity; protein tyrosine phosphatase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA