# Recombinant Human PTK2B 293 Cell Lysate

**Cat. No.** PTK2B-2698HCL **Lot. No.** (See product label)

## SPECIFICATION

| | |
|---|---|
| **Product Overview** | |
| **Species** | Human |
| **Source** | HEK293 |
| **Description** | Antigen standard for PTK2B protein tyrosine kinase 2 beta (PTK2B), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection. |
| **Components** | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| **Size** | 0.1 mg |
| **Storage Instruction** | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment. |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| **Applications** | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane. |

## GENE INFORMATION

| | |
|---|---|
| **Gene Name** | PTK2B PTK2B protein tyrosine kinase 2 beta [ Homo sapiens ] |
| **Official Symbol** | PTK2B |
| **Synonyms** | PTK2B; PTK2B protein tyrosine kinase 2 beta; FAK2, protein tyrosine kinase 2 beta; protein-tyrosine kinase 2-beta; CADTK; CAKB; PTK; PYK2; RAFTK; FADK 2; CAK-beta; protein kinase B; focal adhesion kinase 2; cell adhesion kinase beta; proline-rich tyrosine kinase 2; calcium-dependent tyrosine kinase; related adhesion focal tyrosine kinase; calcium-regulated non-receptor proline-rich tyrosine kinase; PKB; FAK2; FADK2; |
| **Gene ID** | 2185 |
| **mRNA Refseq** | NM_173174 |
| **Protein Refseq** | NP_775266 |
| **MIM** | 601212 |
| **UniProt ID** | Q14289 |
| **Chromosome Location** | 8p21.1 |
| **Pathway** | Alpha-synuclein signaling, organism-specific biosystem; B Cell Receptor Signaling |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| | Pathway, organism-specific biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Calcium signaling pathway, organism-specific biosystem; Calcium signaling pathway, conserved biosystem; Cell-Cell communication, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; |
| **Function** | 3-phosphoinositide-dependent protein kinase binding; ATP binding; binding; non-membrane spanning protein tyrosine kinase activity; nucleotide binding; protein binding; protein tyrosine kinase activity; protein tyrosine kinase activity; signal transducer activity; |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA