

Recombinant Human PSME3 293 Cell Lysate

Cat. No. PSME3-2740HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for proteasome (prosome, macropain) activator subunit 3 (PA28 gamma; Ki) (PSME3), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

PSME3 proteasome (prosome, macropain) activator subunit 3 (PA28 gamma; Ki) [Homo sapiens]

Official Symbol

PSME3

Synonyms

PSME3; proteasome (prosome, macropain) activator subunit 3 (PA28 gamma; Ki); proteasome activator complex subunit 3; Ki; PA28 gamma; PA28G; REG GAMMA; PA28gamma; Ki antigen; Ki nuclear autoantigen; proteasome activator 28-gamma; 11S regulator complex gamma subunit; 11S regulator complex subunit gamma; proteasome activator 28 subunit gamma; activator of multicatalytic protease subunit 3; REG-GAMMA; PA28-gamma;

Gene ID

10197

mRNA Refseq

NM_176863

Protein Refseq

NP_789839

MIM

605129

UniProt ID

P61289

**Chromosome
Location**

17q12-q21

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem;
 APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem;
 APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;
 APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted
 proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C
 and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific
 biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive
 Immune System, organism-specific biosystem;

Function

MDM2 binding; endopeptidase activator activity; p53 binding; protein binding;