# Recombinant Human PSMD7 293 Cell Lysate

**Cat. No.** PSMD7-2745HCL **Lot. No.** (See product label)

## SPECIFICATION

| | |
|---|---|
| **Product Overview** | |
| **Species** | Human |
| **Source** | HEK293 |
| **Description** | Antigen standard for proteasome (prosome, macropain) 26S subunit, non-ATPase, 7 (PSMD7) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection. |
| **Components** | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| **Size** | 0.1 mg |
| **Storage Instruction** | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment. |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| **Applications** | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane. |

## GENE INFORMATION

| | |
|---|---|
| **Gene Name** | PSMD7 proteasome (prosome, macropain) 26S subunit, non-ATPase, 7 [ Homo sapiens ] |
| **Official Symbol** | PSMD7 |
| **Synonyms** | PSMD7; proteasome (prosome, macropain) 26S subunit, non-ATPase, 7; proteasome (prosome, macropain) 26S subunit, non ATPase, 7 (Mov34 homolog); 26S proteasome non-ATPase regulatory subunit 7; MOV34; Mov34 homolog; P40; Rpn8; S12; proteasome subunit p40; 26S proteasome regulatory subunit S12; 26S proteasome regulatory subunit rpn8; Moloney leukemia virus-34 proviral integration; proteasome (prosome, macropain) 26S subunit, non-ATPase, 7 (Mov34 homolog); |
| **Gene ID** | 5713 |
| **mRNA Refseq** | NM_002811 |
| **Protein Refseq** | NP_002802 |
| **MIM** | 157970 |
| **UniProt ID** | P51665 |
| **Chromosome Location** | 16q22.3 |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| **Pathway** | APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem; APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA