

Recombinant Human PSMD6 293 Cell Lysate

Cat. No. PSMD6-2746HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for proteasome (prosome, macropain) 26S subunit, non-ATPase, 6 (PSMD6) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

PSMD6 proteasome (prosome, macropain) 26S subunit, non-ATPase, 6 [Homo sapiens]

Official Symbol

PSMD6

Synonyms

PSMD6; proteasome (prosome, macropain) 26S subunit, non-ATPase, 6; 26S proteasome non-ATPase regulatory subunit 6; KIAA0107; p44S10; Rpn7; S10; p42A; 26S proteasome regulatory subunit S10; 26S proteasome regulatory subunit RPN7; breast cancer-associated protein SGA-113M; phosphonoformate immuno-associated protein 4; proteasome regulatory particle subunit p44S10; SGA-113M;

Gene ID

9861

mRNA Refseq

NM_014814

Protein Refseq

NP_055629

UniProt ID

Q15008

Chromosome Location

3p14.1

Pathway

APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem;

Function

ATPase activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA