

Recombinant Human PSMC5 293 Cell Lysate

Cat. No. PSMC5-2759HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for proteasome (prosome, macropain) 26S subunit, ATPase, 5 (PSMC5) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [PSMC5 proteasome \(prosome, macropain\) 26S subunit, ATPase, 5 \[Homo sapiens \]](#)

Official Symbol PSMC5

Synonyms

PSMC5; proteasome (prosome, macropain) 26S subunit, ATPase, 5; 26S protease regulatory subunit 8; p45; p45/SUG; S8; SUG 1; SUG1; TBP10; TRIP1; MSUG1 protein; proteasome subunit p45; thyroid receptor interactor 1; Tat-binding protein homolog 10; proteasome 26S ATPase subunit 5; proteasome 26S subunit ATPase 5; 26S proteasome AAA-ATPase subunit RPT6; thyroid hormone receptor-interacting protein 1; SUG-1;

Gene ID [5705](#)

mRNA Refseq [NM_001199163](#)

Protein Refseq [NP_001186092](#)

MIM [601681](#)

UniProt ID [P62195](#)

Chromosome Location 17q23.3

Pathway APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem;
 APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;
 APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted
 proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C
 and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific
 biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive
 Immune System, organism-specific biosystem;

Function

ATP binding; ATPase activity; hydrolase activity; nucleotide binding; protein binding;
 receptor binding; thyrotropin-releasing hormone receptor binding; transcription
 cofactor activity; transcription factor binding;