

Recombinant Human PRDM4 293 Cell Lysate

Cat. No. PRDM4-2885HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for PR domain containing 4 (PRDM4) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

PRDM4 PR domain containing 4 [Homo sapiens]

Official Symbol

PRDM4

Synonyms

PRDM4; PR domain containing 4; PR domain zinc finger protein 4; PFM1; PR domain-containing protein 4; PR-domain zinc-finger protein PFM1; MGC45046;

Gene ID

11108

mRNA Refseq

NM_012406

Protein Refseq

NP_036538

MIM

605780

UniProt ID

Q9UKN5

**Chromosome
Location**

12q23-q24.1

Pathway

Neurotrophin signaling pathway, organism-specific biosystem; Neurotrophin signaling pathway, conserved biosystem; Signal Transduction, organism-specific biosystem; Signalling by NGF, organism-specific biosystem; p75 NTR receptor-mediated signalling, organism-specific biosystem; p75(NTR)-mediated signaling, organism-specific biosystem; p75NTR negatively regulates cell cycle via SC1, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem;

Function

DNA binding; death receptor binding; metal ion binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA