

Recombinant Human PPM1G 293 Cell Lysate

Cat. No. PPM1G-2959HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for protein phosphatase 1G (formerly 2C), magnesium-dependent, gamma isoform (PPM1G), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

PPM1G protein phosphatase, Mg²⁺/Mn²⁺ dependent, 1G [Homo sapiens]

Official Symbol

PPM1G

Synonyms

PPM1G; protein phosphatase, Mg²⁺/Mn²⁺ dependent, 1G; protein phosphatase 1G (formerly 2C), magnesium dependent, gamma isoform; protein phosphatase 1G; PP2C; gamma; PP2CG; PP2Cgamma; protein phosphatase 2C; gamma isoform; PP2C-gamma; PP2C, gamma; protein phosphatase 1C; protein phosphatase 2C gamma isoform; protein phosphatase 2C isoform gamma; protein phosphatase magnesium-dependent 1 gamma; protein phosphatase 2, catalytic subunit, gamma isoform; protein phosphatase 1G (formerly 2C), magnesium-dependent, gamma isoform; PPP2CG; PP2CGAMMA; MGC1675; MGC2870;

Gene ID

5496

mRNA Refseq

NM_002707

Protein Refseq

NP_002698

MIM

605119

UniProt ID

O15355

**Chromosome
Location**

2p23.3

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	mRNA processing, organism-specific biosystem;
Function	catalytic activity; hydrolase activity; metal ion binding; protein serine/threonine phosphatase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA