

Recombinant Human POLR2C 293 Cell Lysate

Cat. No. POLR2C-3036HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for polymerase (RNA) II (DNA directed) polypeptide C, 33kDa (POLR2C) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name POLR2C polymerase (RNA) II (DNA directed) polypeptide C, 33kDa [Homo sapiens]

Official Symbol POLR2C

Synonyms POLR2C; polymerase (RNA) II (DNA directed) polypeptide C, 33kDa; polymerase (RNA) II (DNA directed) polypeptide C (33kD); DNA-directed RNA polymerase II subunit RPB3; RNA polymerase II subunit 3; RPB3; RPB33; RNA polymerase II subunit B3; DNA-directed RNA polymerase II subunit C; DNA-directed RNA polymerase II 33 kDa polypeptide; RPB31; hRPB33; hsRPB3;

Gene ID 5432

mRNA Refseq NM_032940

Protein Refseq NP_116558

MIM 180663

UniProt ID P19387

Chromosome Location 16q13-q21

Pathway Abortive elongation of HIV-1 transcript in the absence of Tat, organism-specific biosystem; DNA Repair, organism-specific biosystem; Disease, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Dual incision reaction in TC-NER, organism-specific biosystem; Eukaryotic Transcription Initiation, organism-specific biosystem; Formation of HIV-1 elongation complex containing HIV-1 Tat, organism-specific biosystem; Formation of HIV-1 elongation complex in the absence of HIV-1 Tat, organism-specific biosystem;

Function

DNA binding; DNA-directed RNA polymerase activity; protein dimerization activity; contributes_to protein kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA