

Recombinant Human PLCB1 293 Cell Lysate

Cat. No. PLCB1-3131HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for phospholipase C, beta 1 (phosphoinositide-specific) (PLCB1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [PLCB1 phospholipase C, beta 1 \(phosphoinositide-specific\) \[Homo sapiens \]](#)

Official Symbol PLCB1

Synonyms PLCB1; phospholipase C, beta 1 (phosphoinositide-specific); 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase beta-1; KIAA0581; PLC I; PLC154; PLC-beta-1; phospholipase C-I; phosphoinositidase C; phospholipase C-beta-1; inositoltrisphosphohydrolase; triphosphoinositide phosphodiesterase; phosphoinositide phospholipase C-beta-1; monophosphatidylinositol phosphodiesterase; 1-phosphatidyl-D-myo-inositol-4,5-bisphosphate; 1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase beta 1; 1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase beta-1; PLC-I; EIEE12; PI-PLC; PLCB1A; PLCB1B; PLC-154; FLJ45792;

Gene ID [23236](#)

mRNA Refseq [NM_015192](#)

Protein Refseq [NP_056007](#)

MIM [607120](#)

UniProt ID [Q9NQ66](#)

Chromosome 20p12

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Activation of Kainate Receptors upon glutamate binding, organism-specific biosystem; African trypanosomiasis, organism-specific biosystem; African trypanosomiasis, conserved biosystem; Alzheimers disease, organism-specific biosystem; Alzheimers disease, conserved biosystem; Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem;
Function	GTPase activator activity; calcium ion binding; calmodulin binding; enzyme binding; hydrolase activity; lamin binding; phosphatidylinositol phospholipase C activity; phosphatidylinositol phospholipase C activity; phosphatidylinositol phospholipase C activity; phosphatidylinositol phospholipase C activity; phosphatidylinositol-4,5-bisphosphate binding; phospholipase C activity; protein homodimerization activity; signal transducer activity;