

Recombinant Human PGM3 293 Cell Lysate

Cat. No. PGM3-3249HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for phosphoglucomutase 3 (PGM3) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

PGM3 phosphoglucomutase 3 [Homo sapiens]

Official Symbol

PGM3

Synonyms

PGM3; phosphoglucomutase 3; phosphoacetylglucosamine mutase; acetylglucosamine phosphomutase; AGM1; DKFZP434B187; PAGM; N-acetylglucosamine-phosphate mutase 1; PGM 3; FLJ11614; FLJ13623; DKFZp434B187;

Gene ID

5238

mRNA Refseq

NM_001199917

Protein Refseq

NP_001186846

MIM

172100

UniProt ID

O95394

Chromosome Location

6q14.1-q15

Pathway

Amino sugar and nucleotide sugar metabolism, organism-specific biosystem; Amino sugar and nucleotide sugar metabolism, conserved biosystem; Asparagine N-linked glycosylation, organism-specific biosystem; Biosynthesis of the N-glycan precursor

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



(dolichol lipid-linked oligosaccharide, LLO) and transfer to a nascent protein, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; Post-translational protein modification, organism-specific biosystem; Synthesis of UDP-N-acetyl-glucosamine, organism-specific biosystem;

Function

isomerase activity; magnesium ion binding; phosphoacetylglucosamine mutase activity; phosphoglucomutase activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA