

# Recombinant Human PCCA 293 Cell Lysate

**Cat. No.** PCCA-3400HCL    **Lot. No.** (See product label)

## SPECIFICATION

### Product Overview

**Species** Human

**Source** HEK293

### Description

Antigen standard for propionyl Coenzyme A carboxylase, alpha polypeptide (PCCA), nuclear gene encoding mitochondrial protein, transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

### Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

**Size** 0.1 mg

### Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Applications**

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

**GENE INFORMATION**
**Gene Name**

PCCA propionyl CoA carboxylase, alpha polypeptide [ Homo sapiens ]

**Official Symbol**

PCCA

**Synonyms**

PCCA; propionyl CoA carboxylase, alpha polypeptide; propionyl Coenzyme A carboxylase, alpha polypeptide; propionyl-CoA carboxylase alpha chain, mitochondrial; PCCase alpha subunit; pccA complementation group; propanoyl-CoA:carbon dioxide ligase alpha subunit;

**Gene ID**

5095

**mRNA Refseq**

NM\_000282

**Protein Refseq**

NP\_000273

**MIM**

232000

**UniProt ID**

P05165

**Chromosome Location**

13q32

**Pathway**

2-oxobutanoate degradation I, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Glyoxylate and dicarboxylate metabolism, organism-specific biosystem; Glyoxylate and dicarboxylate

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

metabolism, conserved biosystem; Metabolic pathways, organism-specific biosystem;  
Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins,  
organism-specific biosystem;

**Function**

ATP binding; biotin binding; biotin carboxylase activity; enzyme binding; ligase  
activity; metal ion binding; nucleotide binding; propionyl-CoA carboxylase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA