

Recombinant Human P2RX5 293 Cell Lysate

Cat. No. P2RX5-3498HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for purinergic receptor P2X, ligand-gated ion channel, 5 (P2RX5), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

P2RX5 purinergic receptor P2X, ligand-gated ion channel, 5 [Homo sapiens]

Official Symbol

P2RX5

Synonyms

P2RX5; purinergic receptor P2X, ligand-gated ion channel, 5; P2X purinoceptor 5; P2X5; ATP receptor subunit; ionotropic ATP receptor P2X5; lymphoid-restricted histocompatibility antigen-1; LRH-1; P2X5R; MGC47755;

Gene ID

5026

mRNA Refseq

NM_002561

Protein Refseq

NP_002552

MIM

602836

UniProt ID

Q93086

**Chromosome
Location**

17p13.3

Pathway

Calcium signaling pathway, organism-specific biosystem; Calcium signaling pathway, conserved biosystem; Neuroactive ligand-receptor interaction, organism-specific biosystem; Neuroactive ligand-receptor interaction, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

ATP binding; extracellular ATP-gated cation channel activity; extracellular ATP-gated cation channel activity; ion channel activity; purinergic nucleotide receptor activity; purinergic nucleotide receptor activity; receptor activity; transmembrane signaling receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA