

Recombinant Human OXCT1 293 Cell Lysate

Cat. No. OXCT1-3508HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for 3-oxoacid CoA transferase 1 (OXCT1), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[OXCT1 3-oxoacid CoA transferase 1 \[Homo sapiens \]](#)

Official Symbol

OXCT1

Synonyms

OXCT1; 3-oxoacid CoA transferase 1; 3 oxoacid CoA transferase , OXCT; succinyl-CoA:3-ketoacid-coenzyme A transferase 1, mitochondrial; SCOT; SCOT-s; 3-oxoacid-CoA transferase 1; succinyl CoA:3-oxoacid CoA transferase; succinyl-CoA:3-ketoacid-CoA transferase; somatic-type succinyl CoA:3-oxoacid CoA-transferase; somatic-type succinyl-CoA:3-oxoacid-CoA-transferase; OXCT;

Gene ID

[5019](#)

mRNA Refseq

[NM_000436](#)

Protein Refseq

[NP_000427](#)

MIM

[601424](#)

UniProt ID

[P55809](#)

**Chromosome
Location**

5p13

Pathway

Butanoate metabolism, organism-specific biosystem; Butanoate metabolism, conserved biosystem; Fatty acid, triacylglycerol, and ketone body metabolism,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem; Ketone body metabolism, organism-specific biosystem;
Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins,
organism-specific biosystem; Synthesis and Degradation of Ketone Bodies,
organism-specific biosystem;

Function

3-oxoacid CoA-transferase activity; 3-oxoacid CoA-transferase activity; protein
homodimerization activity; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA