

Recombinant Human NUDT2 293 Cell Lysate

Cat. No. NUDT2-3648HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for nudix (nucleoside diphosphate linked moiety X)-type motif 2 (NUDT2), transcript variant 3 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [NUDT2 nudix \(nucleoside diphosphate linked moiety X\)-type motif 2 \[Homo sapiens \]](#)

Official Symbol NUDT2

Synonyms

NUDT2; nudix (nucleoside diphosphate linked moiety X)-type motif 2; APAH1; bis(5-nucleosyl)-tetraphosphatase [asymmetrical]; Ap4A hydrolase 1; Ap4Aase; bis(5 nucleosyl) tetraphosphatase (asymmetrical); diadenosine 5, 5 P1; P4 tetraphosphate pyrophosphohydrolase; diadenosine tetraphosphatase; nudix motif 2; nucleoside diphosphate-linked moiety X motif 2; bis(5-nucleosyl)-tetraphosphatase (asymmetrical); diadenosine 5,5-P1,P4-tetraphosphate pyrophosphohydrolase; diadenosine 5,5-P1,P4-tetraphosphate asymmetrical hydrolase; MGC10404;

Gene ID [318](#)

mRNA Refseq [NM_147173](#)

Protein Refseq [NP_671702](#)

MIM [602852](#)

UniProt ID [P50583](#)

Chromosome Location 9p13

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Purine metabolism, organism-specific biosystem; Purine metabolism, conserved biosystem; Pyrimidine metabolism, organism-specific biosystem; Pyrimidine metabolism, conserved biosystem;
Function	GTP binding; bis(5-nucleosyl)-tetraphosphatase (asymmetrical) activity; bis(5-nucleosyl)-tetraphosphatase (symmetrical) activity; hydrolase activity; nucleotide binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA